

SLIPPER ORCHIDS

JOURNAL OF

The Slipper Orchid Alliance



Paphiopedilum canhii



Paphiopedilum rungsuriyanum

IN THIS ISSUE:

FIELD SURVEY OF PAPHIOPEDILUM CANHII: FROM DISCOVERY TO EXTINCTION

By Leonid V. Averyanov, Pham Van The, Loc Phan Ke, Hiep Nguyen Tien, Canh Chu Xuan, Vinh Nguyen Tien, and Hieu Nguyen Quang pg 2

PAPHIOPEDILUM RUNGSURIYANUM - A NEW SPECIES DISCOVERED IN NORTHERN LAOS

By Olaf Grusspg 12

21st WORLD ORCHID CONFERENCE pg 13

INTERNATIONAL SLIPPER SYMPOSIUM AND AOS FALL MEMBERS' MEETING..... pg 13

59th PAPHIOPEDILUM GUILD and 1st WORLD SLIPPER ORCHID CONFERENCEpg 14

NEW BOOK ON LATIN-AMERICAN LADY-SLIPPERS

By Olaf Grusspg 15

UPCOMING EVENTSpg 15

FIELD SURVEY OF *PAPHIOPEDILUM CANHII*: FROM DISCOVERY TO EXTINCTION

By Leonid V. Averyanov, Pham Van The, Loc Phan Ke, Hiep Nguyen Tien,
Canh Chu Xuan, Vinh Nguyen Tien, and Hieu Nguyen Quang

Adapted for *Slipper Orchids* by Sandy Öhlund

Introduction

There are events that happen slowly and some that occur so quickly that no one is aware of the danger: extinction of flora and fauna is imminent in many unexplored areas of the world. It has become much easier to change the landscape and accelerate the extinction of species that inhabit primary forests and untouched habitats. Logging, mining and agriculture take their toll. But the photo of a new species on social media will immediately bring collectors, both local and foreign, who invade these pristine areas and decimate the habitat in hopes of a quick profit. Because of this, the recently discovered *Paphiopedilum canhii* is now an endangered species with a precarious future. And there we have it: extinction at the speed of the Internet.

Botanists and biologists painstakingly chart the situation, but their field studies are filed away in libraries, herbariums and dusty natural history museums; those who might help the situation are not made aware of the impending catastrophes and extinctions. Some interested parties choose to make solutions more difficult with bureaucratic rules and laws that are counterproductive to saving the habitats and species in peril.

Here we have adapted a field investigation on the situation of *Paphiopedilum canhii* by Leonid Averyanov and his group, in order to make their findings more available. We are grateful to have field surveys to make the public aware of the tenuous existence of species such as *P. canhii*, and to encourage meaningful action to save critically endangered endemic species. We are indebted to Prof. Averyanov and his colleagues for their work and acknowledge the sponsors of this important study. (For additional information and references, please see “*Paphiopedilum canhii* in Laos Phou Phachao Mountain – Mountain of *Paphiopedilum canhii*” by Averyanov, *et al.*, in *Orchid Digest*, 78:2, pp. 96-101.)

Sandy Öhlund

Summary

A research expedition program funded by The Rufford Small Grant Foundation, the American Orchid Society and the Chicago Zoological Society (Chicago Board of Trade Endangered Species Fund) to undertake a field assessment of the recently described species *Paphiopedilum canhii* has been completed. New original data relating to habitat and natural conditions of five subpopulations have been obtained. Associated plant species and vegetation have been described and documented. The factors leading to rapid extinction are analyzed and discussed. The data obtained support an assessment of the present status of the species as critically endangered, approaching full extinction in the near future. Conservation of this species in its natural habitats (Fig. 1) would require very urgent actions that may be unrealistic, and *ex-situ* cultivation of the species may be the only way to effectively prevent its imminent extinction. Some recommendations for approaches to orchid protection are briefly proposed.



Fig. 1. *Paphiopedilum canhii* in its natural habitat on a vertical limestone cliff

Background

The rocky limestone formations of northern Vietnam contain gravely endangered primary forest that harbors remnants of a unique area of *Paphiopedilum* speciation and diversity. Studies of this area have revealed the existence of more than 25 isolated endemic species of *Paphiopedilum*. Many of these species have been described by

authors outside of Vietnam; very little information has been available about these species, and the natural habitats are often destroyed by the time the new species are described. Field studies are undertaken, but over-collecting has often decimated the population of the new taxa. Local endemic species such as *P. vietnamense* and *P. tranlienianum* have become extinct by the time of their description. In addition to commercial collecting, deforestation and total habitat collapse have led the way to species extinction.

For the recently described *Paphiopedilum canhii*, the Vietnamese NGO The Center for Plant Conservation (Vietnam Union of Science and Technology Associations) organized field investigations. Under the guidance of Center director, Dr. Nguyen Tien Hiep, and the Vice-Dean of the Faculty of Science (National University of Laos), Dr. Somchanh Bounphanmy, the main goals were delimiting the distribution of the species, describing the habitats, determining the conservation status with International Union for Conservation of Nature (IUCN) criteria, and suggesting protection strategies.

When *Paphiopedilum canhii* arrived in the local markets of Dien Bien and Son La, there was no information on its provenance. Since the limestone regions of these provinces along the Laotian border are known to be the home of strictly endemic orchid species such as *Bulbophyllum paraemarginatum*, *Dendrobium farinatum*, *D. trantuanii*, *D. vietnamense*, *Hayata glandulifera*, *Paphiopedilum x asperum*, *P. barbigerum* var. *barbigerum* (*coccineum*), and *Sunipia nigricans*, Averyanov and his colleagues predicted that *P. canhii* had a restricted, very limited habitat in these rocky, karstic limestone regions of northwestern Vietnam and northeastern Laos. (Fig. 2)



Fig. 2
Area of field explorations for *Paph. canhii* in Indochina and Vietnam

History of the Discovery of *Paphiopedilum canhii*

The Vietnamese people like orchids. Orchid collections and family orchid gardens abound in Vietnam. Orchids for home cultivation can be easily purchased in local neighborhood markets or simply along roads where they pass through areas with remnants of primary forests; these plants are usually offered for sale by local people. They commonly gather these plants from nearby areas, within degraded primary forest, or while forests are being either logged or burned for agricultural fields. These fields permanently replace the primary forests in botanically unexplored areas. Orchid collecting in these places sometimes results in the discovery of rare or undescribed orchids. Such is the story of the discovery of *Paphiopedilum canhii*.

In fact, *Paphiopedilum canhii* was discovered at the end of 2009 by unknown H'Mong (Meo) people who brought plants from a remote mountainous area to sell in Dien Bien City. A few of these plants came into the hands of Mr. Chu Xuan Canh, who cared for them in his garden; they flowered for him the following March. Soon it was evident that these plants were an undescribed species, and the taxon was named *Paphiopedilum canhii*, in honor of the man who flowered the cultivated plants. The new species was published by Prof. Averyanov in the May, 2010 issue of *Orchids*, the journal of the American Orchid Society. Due to a technical error, the species was republished later in Prof. Averyanov's monograph – "*The Orchids of Vietnam, Illustrated Survey, Part 2, Subfamily Orchidoideae*" in the second-quarter issue of volume 13 of the Siberian botanical magazine – *Turczaninowia*.

Publication of a newly discovered species attracts a great deal of attention from people involved in orchidology and horticulture. It also generates interest within the domestic and international orchid trade. The increased market demand for a newly described species leads to a rapid price escalation, which in turn stimulates plant collecting in the area. As a result, large numbers of plants of *P. canhii* immediately appeared in the local markets. In addition, the species' native habitat, long unknown, was explored by dealers and local collectors. Since it was determined that *P. canhii* was an uncommon endemic species with a very restricted habitat, an investigation of the area was mounted in order to gather information before the new species became extinct.

Methodology and Approaches

In planning the investigation, information was gathered from many sources: the local orchid markets, orchid enthusiasts, growers, and dealers in the area of Dien Bien

and Son La provinces. Additional data were obtained from local people, village authorities and local foresters. This information led to the conclusion that *Paphiopedilum canhii* likely occurred in the area of Dien Bien and Son La provinces near the Laotian border, most probably on rocky limestone. Since all slipper orchids grow in intact primary forests, any disruption or destruction of those forests would limit the areas of possible field research. The field overview and special surveys indicate that more than 99.5% of primary forest is gone. The rare remnant forests were found in just a few remote mountain regions.

Landscape and Landforms in the Area of *Paphiopedilum canhii* Habitats

Paphiopedilum canhii is known from a single locality of Dien Bien province near the Laotian border (Fig. 3). This area contains alluvial valleys, shale hills and low mountains, with isolated, rocky mesas or mountains of highly eroded limestone (Fig. 4). These are spectacular formations with rocky, vertical cliffs commonly up to 2600 to 3250 ft above sea level (800-1000 m), but occasionally reaching as high as 4200 to 4600 ft a.s.l. (1300-1400 m) (Fig. 5). Such habitats are home to many native slipper orchids. Most inaccessible mountain tops or cliffs in northwestern Vietnam still retain remnants of rich primary forests. The highest level of species diversity and endemism was observed in this area compared to any other mountain formations in the country. During the fieldwork in these degraded remnant forest patches, numerous rare and strictly endemic plant species were located, including *Paphiopedilum canhii*.



Fig. 3. Studied area of NE Indochina (outlined with red line) and localities of field work designated with black figures.




Fig. 4: Isolated massifs of rocky limestone hills and mountains in *Paphiopedilum canhii* area.



Fig. 5. Typical remnant eroded limestone mountains in Son La Province at elevations of 2600-3250 ft.

A small number of subpopulations inhabit isolated massifs of only a few scattered remnant limestone hills. These areas are highly eroded, marble-like rocky limestone with numerous steep slopes and vertical cliffs. These limestone formations topographically occur between rolling shale hills on the watershed of western tributaries of the Nam He River, but some of them tower above flat river valleys. The total area of this hilly limestone massif does not exceed 6.25 sq miles (4 km²).

The tops of remnant limestone hills at 3000 to 3900 ft a.s.l. (950-1200 m) still support severely fragmented, degraded pieces of primary forest that provide home to native species. These limestone refuges are surrounded by shale hills covered with weedy, secondary plant communities and valleys occupied by agricultural fields. Nine small primary forest stands were located in the area of *Paphiopedilum canhii* populations that might support acceptable habitats, but *P. canhii* populations were found in only five of these sites and in every case as very small subpopulations.

Main Vegetation in the Area of Study

Rich, primary, broad-leaved evergreen, closed submontane tropical forests represent the typical original pristine vegetation type within the area of *Paphiopedilum canhii* (Fig. 6). The presence of very diverse species, specific horizontal and vertical forest structures, permanent shade and humidity, as well as an absence of exotic or weed



Fig. 6: Rich, primary, broad-leaved evergreen, humid tropical forests in the *Paphiopedilum canhii* area

species are typical features of this native vegetation type. Unfortunately, the primary forest has largely been replaced by degraded secondary forest and scrub vegetation with many introduced weedy species.

In the habitat of *Paphiopedilum canhii*, there are many evergreen trees interspersed with a high number of deciduous trees, especially on mountain tops, south-facing slopes and cliffs. In spring these trees produce flowering canopies as the young leaves are developing.

Many weedy species appear quickly in fragments of primary woods opened by forest fires or logging. In the habitats of *P. canhii* there are numerous lithophytic and epiphytic plant species, including many ferns and orchids, as well as lianas, creeping and climbing species, mosses and lichens, often forming mats covering the rocky outcrops on the tops of hills and mountains. Epiphytes often completely cover the branches of old, gnarled trees.

Characteristics of *Paphiopedilum canhii* Habitats and Common Lithophytic Associates

Shady vertical cliffs with northern, northeastern and northwestern exposures on steep rocky slopes of remnant limestone hills and mountains covered with intact primary broad-leaved evergreen forest were consistent features of the habitat of all the subpopulations of *Paphiopedilum canhii*. The species grows on the upper portion of hilly slopes and high vertical cliffs, usually between 130 to 200 ft (40-60 m) below the hilltops (Fig. 7). However, occasionally plants almost reach the summits of these mountains where they grow in shady humid crevices. Plants typically grow as lithophytes on rough solid vertical walls covered with scattered mosses in constant humidity and in permanent shade of the canopy of primary forest. Very often the plants' roots are attached only to solid unbroken limestone walls without any fissures or holes. The remnant primary forests supporting all populations of *P. canhii* studied were still intact but very endangered, given the constant threat of logging.

Numerous lithophytic ferns and orchids are commonly found growing with *P. canhii* in its native habitats. Orchids in the same area include several *Bulbophyllums*, *Calanthe triplicata*, *Coelogyne fimbriata*, *Dendrobium chrysanthum*, several *Epigeniums*, *Liparis*, *Oberonias*, and others. Furthermore, *P. canhii* sometimes grows with other slipper orchids, such as *P. concolor* and *P. dianthum* and in close proximity to populations of *P. barbigerum* var. *coccineum* and *P. malipoense*, that theoretically provide the possibility for the creation of natural hybrids.



Fig. 7: *Paphiopedilum canhii* in its natural habitat, growing on vertical cliffs covered with orchids and mosses, in stable humidity and shade

***Paphiopedilum canhii* in its Native Habitats and Estimation of its Present Status**

In this investigation, *Paphiopedilum canhii* was found in five subpopulations on isolated limestone mountains in an area of less than 0.19 mi² (0.5 km²). The total area of probable species occurrence is even smaller, estimated at an area of 0.135 mi² (0.35 km²). In fact, the actual area of this species growing on vertical cliffs is certainly less than a few hundred square meters. The total number of specimens in all five subpopulations before commercial collecting may be estimated at between 10,000-15,000 mature (flowering-size) individuals, and it is possible that these five subpopulations represent the total extant area of *P. canhii*. In December of 2010, no intact colony was located and the majority of colonies in all five subpopulations had been completely collected. Only occasionally, on the most inaccessible cliffs, were remnants found of poor quality juvenile or weak, stressed specimens. Decimation of previously larger colonies was 99 to 100%. It is estimated that between 2009 and 2010, 99% of the plants in the five studied subpopulations were removed by collectors, and that surviving mature (flowering size) specimens in all subpopulations represent 0.01% of the species' former population (see Fig. 7). As commercial collecting will not immediately cease, these are probably the last images of *P. canhii* in nature.

Based on these data and current original observations, *Paphiopedilum canhii* is presently estimated in IUCN terms as a critically endangered species, reflecting the fact that this species faces an extremely high risk of imminent extinction in the wild.

Taxonomy, Morphology and Biology of *Paphiopedilum canhii*

Paphiopedilum canhii was described by Prof. Leonid Averyanov and Mr. Olaf Gruss in 2010 on the basis of a few specimens of uncertain origin, purportedly collected in Son La Province and flowered in March of 2010 in Hanoi in a private orchid collection belonging to Mr. Chu Xuan Canh.

Later, the project participants found the species in Dien Bien Province. Extensive broad searches for *P. canhii* outside this area in Vietnam and Laos have yielded no positive results. Certainly, this species should now be regarded as a local Vietnamese endemic with very restricted distribution in the province of Dien Bien, northwestern Vietnam.

For a complete taxonomical description of *Paphiopedilum canhii*, please see the original description in:

Leonid V. Averyanov, Olaf Gruss, Canh Chu Xuan, Loc Phan Ke, Dang Bui, Hiep Nguyen Tien. 2010. *Paphiopedilum canhii*. A New Species from Northern Vietnam. *Orchids*, Vol. 79, No 5, pp. 288-290.

History of the Exploitation and Extinction of *Paphiopedilum canhii*

Extensive exploitation of *Paphiopedilum canhii* started immediately after its discovery in mid-2009. The species was first found by members of the local H'Mong (Meo) minority and collected along with other orchids for sale in the local markets of Dien Bien and Son La cities. This very distinct and unusual slipper orchid was highly valued by local orchid lovers, and in just a few weeks, rumors of an intriguing new species spread widely, reaching Hanoi and other large cities of northern Vietnam. The first specimens of this mysterious plant flowered at the beginning of 2010 in a number of private collections in Hanoi, Dien Bien and Son La cities. In the next few days, the internet was filled with images of this new flower. Illustrated publication of this new species in the world-renowned American Orchid Society *Orchids* magazine stimulated market demand and supported a rapid price escalation. Numerous international dealers arrived in Son La and Dien Bien cities in the spring of 2010; their purpose was to hunt for and trade this plant. The best clones at this time were traded at a cost of up to US \$100 for a single plant. The largest shipments of plants were supposedly imported into Taiwan and Europe.

Exciting rumors about high prices for plants provoked true orchid fever in the habitat of *Paphiopedilum canhii*. Local people from neighboring villages put aside their daily duties and, instead, went to the forest; on some days, more than 20 plant hunters collected plants.

Naturally, this large influx of plants rapidly exceeded the demand and very soon prices fell from US \$100 per plant to US \$50 per kilogram. By the end of 2010, prices had continued to fall and prices of US \$10-20 per kilo were common. Even at these prices, trade had seriously weakened; nearly all plants that did not sell were simply destroyed. Ironically, the wholesale collecting of the great majority of plants from their habitat was for no purpose. They brought no money to local people, no profits to local or international traders and no happiness to orchid lovers all over the world. These last specimens of this unique, critically endangered species, threatened with imminent extinction, were simply wasted. The lack of horticultural knowledge and experience necessary for successful slipper orchid cultivation prevents *P. canhii* from being successfully grown in local collections and nurseries. Even large flowering-size plants have little chance of survival in these conditions (Fig. 8). At the end of 2010, trade in *P. canhii* had completely stopped for three main reasons: having already purchased the best cultivars necessary for propagation and breeding, demand from foreign traders was gone, due to the difficulties encountered



Fig. 8: *Paphiopedilum canhii* in local nurseries of Dien Bien City

in cultivation; demand from domestic purchasers was gone; and the very few left in nature had made collecting them unprofitable.

As a very approximate estimate, 25-35 kg of *P. canhii* plants were collected during the short history of its discovery, exploitation and extinction in the wild. Since they are fairly small plants, 1 kg (2.2 pounds) of freshly-collected material contains about 300-350 mature (flowering-size) plants and commonly 200-300 juveniles. This suggests that at least 10,000-15,000 mature and juvenile specimens were collected. Direct observation in nature indicates that about 99.5% of the population was extinguished in only six months of exploitation. Cultivation under appropriate conditions, in the best-equipped nurseries, may prevent full species extinction. However, such activity is illegal due to CITES regulations. Not a single CITES export permit application for *P. canhii* has ever been processed, and all specimens of the species outside Vietnam came from the black market and remain illegal.

Remarkable Plant Species Associated with Habitats of *Paphiopedilum canhii*

Remnants of primary forests in the area of *P. canhii* often retain elements of the diverse, highly endemic flora. Many of these species vanished from other regions of Vietnam years ago, but in the studied area, they still survive and sometimes are represented by fairly large populations. The rarest and most interesting species observed here were slipper orchids, such as *P. concolor*, *P. malipoense*, *P. barbigerum* var. *coccineum* and *P. dianthum*. All of these discoveries essentially expand our knowledge about the distribution of these rare slipper orchid species and represent new records for northwestern Vietnam.

About 112 orchid species were observed as direct associates in habitats of *P. canhii* subpopulations. Some very rare orchids listed from a single collection were also found during our field exploration of the *P. canhii* habitat. Among the taxa discovered and proposed for future description as new to science are nine orchid species: a new genus and species - *Lockia sonii* and eight species of the genera *Bulbophyllum*, *Cleisostoma*, *Dendrobium*, *Hippeophyllum*, *Saccolabiopsis*, *Sarcoglyphis* and *Schoenorchis*. Also discovered and proposed for description is a remarkable new species of *Begonia* - *Begonia viscosa* - an important undershrub-dominant of rocky limestone areas of northern Laos.

Widely cultivated as highly prized ornamental plants - *Dendrobium trantuanii* and *Stereochilus brevirachis* - were found in their natural habitats. Both appear as notable

associates of *Paphiopedilum canhii* as does *Vanda brunnea*, a very important ornamental species, abundant in the area studied and with an outstanding variation in flower shape and color.

Main Factors of Orchid Extinction

Deforestation is the leading factor in the extinction of native floras globally. In tropical areas, forest removal leads very rapidly to degradation of the fertility of the soil. The loss of topsoil opens the way to rapid degradation of vegetation and terminates in a climax community of secondary shrub and herbaceous weed species. This process is irreversible. Primary forests with their very sensitive native species never regenerate. Extinction of such forests leads to a catastrophic decrease in the genetic diversity of the world. Such a stark scenario was observed within the area studied.

Primitive slash-and-burn agriculture and forest logging for domestic purposes and fuel are the main reasons for deforestation in areas of rapidly increasing human populations. Degraded pieces of remaining primary forests are presently estimated as less than 0.5% of the territory. Such unique remnant forest areas still occur in a few remote mountain regions. Their protection and conservation are impossible without urgent action of municipal, provincial, and governmental authorities. Without urgent and effective



Fig. 10. Primitive maize fields replacing natural habitat of *Paphiopedilum canhii*



Fig. 9. Logging of canopy of primary forest in *Paphiopedilum canhii* area



Fig. 11. Full collapse of rich, native vegetation through forest burning for agriculture



Fig. 12. Destruction of vegetation and mountains by mining and road construction in Dien Bien Province

action, the forest here will be completely destroyed in the near future. The survival of sensitive native species, including *P. canhii*, after forest destruction will be impossible (Figs. 9, 10, 11, 12).

Construction of roads, highways, mines, high-voltage power and communication lines, as well as any other large land exploitation projects are also important factors in the destruction of nature that follows primary forest extinction. For example, road construction in the area of *P. canhii* destroys not only vegetation, but also even the physical landscape.

Landscape desertification and aggressive weed introgression play the role of slow, but irreversible destructive factors in these last isolated refuges of primary forests in the world's tropical zones and result in higher summer temperatures and loss of air humidity, particularly during the dry seasons. This effect seriously depresses primary humidity-loving vegetation, particularly in its small isolated refuges. Under such conditions, most sensitive species slowly decline even in physically intact primary plant communities. Shade-loving native herbaceous lithophytes growing on vertical cliffs in conditions of permanent humidity represent the most sensitive and endangered group of species. Such species always vanish first, because vertical cliffs lose humidity much faster than other substrates during the

process of habitat destruction. *Paphiopedilum canhii* and its slipper orchid associates belong to just this group of species.

Destruction of native floras by aggressive exotic and weed species represents a huge problem globally. Eventually primary plant communities are transformed into secondary thickets of weed and alien exotic species. This process proceeds without obvious visible cataclysms, but leads to strong depression of native species and eventually kills most of them. This is particularly true for most sensitive and rare endemic orchids, including all species of *Paphiopedilum*.

Commercial collecting is another important factor in the extinction of saleable non-timber plant species. Collection of food, spice and medicinal species is commonplace. However, commercial collecting of ornamental plants often helps to maintain these species in cultivation and may play a positive role in their survival. Collecting of the rarest endemic plants in irreversibly degraded habitats for cultivation under appropriate control may be the only way to conserve unique local plant diversity. This is true for many strictly endemic orchids including *P. canhii*, which could have become completely extinct before its discovery due to deforestation. The uncontrolled commercial collecting of this species has seriously depleted its population, but it has also provided an important chance for the plant's *ex situ* survival. Strictly limited collecting of rare orchids in nature for cultivation and propagation under the control of scientific and state/local authorities in similar situations can provide an interested market with cultivated seedlings and make commercial collecting of plants in nature unprofitable. Such an approach provides a practical means for rare species protection. Any collecting restrictions and prohibitions currently in practice appear to be absolutely ineffective in protecting rare market-valued species.

Summarized Conclusions

Paphiopedilum canhii is endemic to northwestern Vietnam with extremely restricted distribution, located in a single geographical area of less than 0.35 km². Ecologically this species is an obligate lithophyte growing at elevations of 900-1200 m a.s.l. in intact primary forests on cliff faces of remnant mountains composed of highly eroded crystalline rocky limestone.

Paphiopedilum canhii in its native habitats grows in association with a number of extremely rare orchid species that became extinct in other areas of Vietnam many years ago.

Commercial collecting due to international demand was a main factor in the decimation of the *P. canhii* population immediately following its discovery.

The intact total population of the species before mass collection is estimated at approximately 10,000-15,000 mature and juvenile specimens. About 99.5% of this population is presently gone due to uncontrolled commercial collecting.

Progressive deforestation and land exploitation are additional factors that threaten the remaining habitat for the species.

According to IUCN criteria *P. canhii* belongs to a group of critically endangered species approaching full extinction in nature.

Conservation of the species in nature needs immediate effective overall protection of its habitats.

Effective conservation of the species *ex situ* demands legalized collecting and transport of plants into high-technology nurseries.

The example of *P. canhii* distribution confirms the fact that strictly local endemism is very typical for limestone floras of northern Vietnam. Under the conditions of total, uncontrolled deforestation, a large number of similar local endemics will certainly become extinct in this area in the near future, many before they are discovered.

Because among such plants may be species of outstanding economic significance, botanical explorations of this area are urgently necessary for the salvation of global plant diversity and national plant heritage.

National Actions for Protection

Organization and establishment of protected areas for all existing habitats of *P. canhii*, including full conservation of remnant primary forests, strict limitation of land exploitation and effective prohibition of plant collecting are urgently needed. The total area necessary for protection does not exceed 1.54-2.3 mi² (4-6 km²).

Education and training in the design and implementation of conservation actions of local authorities (Municipality People's Committee, Forest Protection Department and associated local agencies) must be done immediately. The realization of an educational program for local people and local authorities with an explanation of the outstanding significance of *P. canhii* conservation for organization of highly-prized ecotourism visitation sites would be a shining example of a locality with the highest orchid diversity in all of the Indochinese Region. In fact, the organization of ecotourism in this area would partially solve the problem of local unemployment.

General Actions for Protection

The immediate creation of a high-technology "rescue center" for the cultivation, propagation and repatriation of critically endangered plant species (including *P. canhii*) is highly recommended. Such a center would be under the oversight of Vietnamese governmental/scientific agencies or in cooperation with official "rescue centers" abroad.

Strict limits on the collecting of rare orchids in nature, as well as cultivation and propagation under control of scientific and state/local authorities in cases similar to *P. canhii*, would fill the market with cultivated seedlings and make the commercial collecting of plants in nature unprofitable. Such approaches provide a real means to rare species protection.

Authorities should facilitate the sale of plants (including orchids) collected from habitats irreversibly damaged by total deforestation. Any restrictions on such sales play a strongly negative role in conservation of world plant diversity.

Immediate improvements to CITES regulations are necessary to minimize the bureaucratic mechanism. It should be recognized by the Parties to the Convention that current measures aimed at plant protection are often ineffective. In its present form, CITES irrationally expends considerable national and international human and financial resources while making no real, objective headway for protection of the natural environment.

Acknowledgements

The fieldwork in Vietnam and Laos was organized by The Center for Plant Conservation (Vietnam Union of Science and Technology Associations) and the National University of Laos, Faculty of Science. Laboratory processing of collected materials was completed at the Komarov Botanical Institute (Russian Academy of Sciences). Expeditions were supported by the following agencies: The Rufford Small Grant Foundation ("Assessment of the Distribution and Natural Status of *Paphiopedilum canhii* in Vietnam"), the American Orchid Society ("Assessment of Orchid Endemism in Northwestern Vietnam with Special Attention to *Paphiopedilum canhii*"), Chicago Zoological Society, Chicago Board of Trade Endangered Species Fund ("Assessment of the Current Natural Status of Critically Endangered Species - *Paphiopedilum canhii* - for Its Conservation"), and the U.S.A. National Geographic Society (grant # 8800-10 - "Exploration of Primary Woods along Constructed Highway Hanoi - Ho Chi Minh for their Sustainable Conservation in Limits of Ha Tinh, Quang

Binh, Quang Tri, Thua Thien Hue, Quang Nam and Kon Tum Provinces of Central Vietnam”). Deep gratitude is due to all these organizations.

Authors

Leonid V. Averyanov, Dr Sci, PhD, is a professor and a vice president of the Russian Botanical Society. He is co-author of *Slipper Orchids of Vietnam*. Komarov Botanical Institute of the Russian Academy of Science, Prof. Popov str. 2, St.-Petersburg, 197376, Russian Federation; E-mail: av_leonid@mail.ru ; av_leonid@yahoo.com.

Pham Van The, M.Sc., is a member of the Institute of Ecology and Biological Resources of Vietnam Academy of Science and Technology; E-mail: phamvthe@gmail.com ; phamvthe@hotmail.com.

Loc Phan Ke, PhD, is a professor at Hanoi University of Science of Vietnam National University, Hanoi. 334 Nguyen Trai, Thanh Xuan, Hanoi, Vietnam, and the Center for Plant Conservation of Vietnam Union of Science and Technology Associations, Lac Long Quan Rd, Nghia Do, Cau Giay, Ha Noi, Vietnam; E-mail: pkeloc@yahoo.com.

Hiep Nguyen Tien, PhD, is director of the Center for Plant Conservation, Vietnam Union of Science and Technology Associations. Lac Long Quan Rd, Nghia Do, Cau Giay, Ha Noi, Vietnam; E-mail: ntienhiiep@hn.vnn.vn.

Canh Chu Xuan, BA, is a natural resources governance field coordination officer with CARE International in Vietnam, 66 Xuan Dieu, Tay Ho, Ha Noi; P101-92A2 Thanh Nhan street, Hai Ba Trung District, Ha Noi, Vietnam; E-mail: cxcanh@gmail.com.

Vinh Nguyen Tien is a staff member of the Center for Plant Conservation, Vietnam Union of Science and Technology Associations. Lac Long Quan Rd, Nghia Do, Cau Giay, Ha Noi, Vietnam; E-mail: ntienhiiep@hn.vnn.vn.

Hieu Nguyen Quang, M.Sc., is vice director of the Center for Plant Conservation, Vietnam Union of Science and Technology Associations. Lac Long Quan Rd, Nghia Do, Cau Giay, Ha Noi, Vietnam; E-mail: hieu.nguyen@cpcvn.org.

Photos by Leonid V. Averyanov & Pham Van The

Adaptor of this Article

Sandy Öhlund is an accredited judge with the American Orchid Society, and specializes in Paphs and Phrags. She is a retired professor, an artist and author; her most recent publication is the 2014 e-book *Paphiopedilum* by Braem, Chiron and Öhlund.

“The Media View”

by Sandy Öhlund



PAPHIOPEDILUM RUNGSURIYANUM
A NEW SPECIES DISCOVERED IN NORTHERN LAOS

by Olaf Gruss

Translated by Judith Rapacz-Hasler

Summary: A new and very distinct species of the genus *Paphiopedilum*, discovered in Northern Laos, is described as *Paphiopedilum rungsuriyanum*.

In recent years a large number of *Paphiopedilums*, as well as other orchids, arrived from Laos to markets in Thailand. Niwat Rungruang, a grower in Thailand, received a group of very small size plants. When they started to bloom in May, 2014, he was surprised that he did not recognize them as a known *Paphiopedilum* species. He contacted orchid specialists in Thailand and Germany to ask for an identification, based on several flowering plants and detailed picture material (supplied by Olaf Gruss from Germany). While the plants with their marbled leaves appeared quite similar to *Paphiopedilum canhii*, the flowers clearly showed much broader petals, an intense red-purple color, as well as an entirely different staminode. Also, the underside of the leaves differed significantly, being gray-green with wide purple veins, in contrast to *P. canhii*, which has red-purple speckles.

This new species, *Paphiopedilum rungsuriyanum*, is named in honor of Niwat Rungruang, using the first part of his name, "Rung," that means flowering period, success, prosperity, and "Suriya," that means growing. Although this new species is closely related to *Paphiopedilum canhii*, it differs by the color of the flower, the shorter and wider petals, the wider lip and the folded, almost ovate staminode. Also, the flower is quite small, with a natural spread of about 2 inches (5.2 cm) horizontal and 2 3/8 inches (6 cm) vertical, compared to 6 and 8 cm, respectively for *Paph. canhii*. The plant is lithophytic or terrestrial with 3-5 opposite leaves, a short stem, and hardly any distance between the individual growths.

This small spectacular species should be extremely valuable to breeders because of its dramatic coloring and compact size that is ideal for window-sill culture. It is hoped that artificial propagation will soon commence so that its natural habitat in Laos can be protected.

LITERATURE

Averyanov L.V., 2008. *The Orchids of Vietnam, Illustrated Survey, Part 1, Subfamilies Apostasioideae, Cypripedioideae and Spiranthoideae*. Turczaninowia, Vol. 11, N 1. pp. 5-168.

Averyanov L.V., Phillip Cribb, Phan Ke Loc, Nguyen Tien Hiep. 2003. *Slipper Orchids of Vietnam, With an Introduction to the Flora of Vietnam*, Royal Botanic Gardens, Kew, Compass Press Limited, 308 p.

Braem, Guido, Guy Chiron & Sandy Öhlund, 2014. *Paphiopedilum e* – book; in English, Moorland, Denmark; ISBN 9788793044012.

Cribb P.J., 1998. *The Genus Paphiopedilum*, Kota Kinabalu, Nat. Hist. Publ. (Borneo), 427 p.

Liu Zhongjian, Chen Sing-chi, P.J. Cribb, 2009. 4. *Paphiopedilum Pfitzer*, pp. 33-44. In: Wu Zhenqyi, P.Raven, Flora of China, Vol. 25, Science Press & MBG Press, Beijing & St. Louis. 570 p.

AUTHORS

Olaf Gruss, In der Au 48; 83224 Grassau; Germany;
 E-mail: a-o.gruss@t-online.de

Niwat Rungrung - 144/348 rangsit-nakonayok44;
 Prachatap Rangsit-nakonayok Rd.; Patumthani
 Thanyaburi; 12130 Thailand

Yongyouth Chaisuriyakul - 14/27 soi soonvijai 6 New
 petchburi Rd.; Bangkok. Hany Kwang; Bangkok 10320
 Thailand

Ibn Dionisio – 6419 Rochdale Boulevard Regina, SK S4X
 2Y9 Canada

The original article about *Paphiopedilum rungsuriyanum*, upon which this article is based, appeared on the website of the *Orchideen Journal* and may be accessed at:

http://orchideenjournal.de/permalink/GRUSS_RUNGRUANG_YONYOUTH_CHAISURIYAKUL_DIONISIO_Paphiopedilum.pdf



Comparison of plants of *Paphiopedilum rungsuriyanum* (left) and *Paphiopedilum canhii* (right) Photo: Olaf Gruss

*Paphiopedilum rungsuriyanum*

Front view of two different clones (left) and a side view (right) Photos: Olaf Gruss



Staminodium with column
Photo: Olaf Gruss

*Paphiopedilum rungsuriyanum*

Measurements: horizontal 2 inches (5.2 cm) and vertical 2 3/8 inches (6 cm)
Photos: Ono Achima

21st WORLD ORCHID CONFERENCE IN SOUTH AFRICA

Slipper lovers who attend the WOC in Johannesburg, September 10-14, 2014, will have the opportunity to meet and socialize at the Slipper Enthusiast Forum on Wednesday, September 10, from 5-7 pm. A panel discussion with growers from around the globe will discuss their experiences with growing and/or hybridizing. It is being organized by Faan Kruger, and if you plan to attend, wish to be part of the panel or have questions, please contact him at info@paphiopedilum.co.za. Faan suggests you wear a Slipper hat to the event and promises a prize "for the best dressed Slipper."

INTERNATIONAL SLIPPER SYMPOSIUM AND AOS FALL MEMBERS' MEETING

The International Slipper Symposium will be held in conjunction with the American Orchid Society Fall Members' Meeting on Oct. 29-Nov. 2, 2014, at the Sheraton Orlando North, Maitland, FL. Saturday, Nov. 1, will be the day to enjoy these Slipper speakers: Manolo Arias on *Phrag. kovachii* culture, Chen-Chung Chen on multifloral paphs in Taiwan and John Doherty on *Paph. sanderianum* (sponsored by the SOA). The SOA will have a meeting and auction following John's lecture on Saturday afternoon at 3:30. Go to the AOS website at www.aos.org and select "Fall Meetings" or the International Slipper Symposium at www.slippersymposium.com for information and registration.



59th Paphiopedilum Guild and 1st World Slipper Orchid Conference

The Orchid Digest is pleased to announce that the 59th Paphiopedilum Guild will be expanded to become the 1st World Slipper Orchid Conference to be held in January 2015 in Hilo, Hawaii.

This enhanced program will include international speakers as well as speakers from the U.S. It is a pleasure to note that the local Hilo orchid members are already geared up for this, holding meetings to ensure that everything runs smoothly. Other groups, including SOA, are getting involved. It should be a weekend to remember. Two full days of lectures, sales, auction, open houses, and great food (and maybe vacation). The venue, Hilo Hawaiian

Hotel, will extend their special room rates for a few days on either side of the conference. Start saving and looking for good buys for airline tickets. If you are an AOS judge, you can receive credit for an out-of-region judging on Friday, January 16. Depending on your judging region, you may also be able to get credit for attending the two days of meetings. You may find more information and register for the meeting by going to the Orchid Digest webpage (www.orchiddigest.org). The costs of the meetings and hotels are very reasonable. Gather up some of your friends and seriously think about joining us.

Orchid Digest

<http://www.castleresorts.com/Home/accommodations/hilo-hawaiian-hotel>

MEMBER ACCESS TO NEWSLETTER ARCHIVE ON THE SOA WEBSITE

username = Newsletter
password = phrag



**New Book on
Latin American Lady-Slippers
(Lateinamerikanische Frauenschuhe)
*Phragmipedium, Mexipedium
Selenipedium***

by Olaf Gruss

Publisher: Orchideenzauber Verlag 2014,
Ruhmannsfelden, Germany.
528 pages, over 2000 color photographs
US \$135 (99.- €), German Text

The book can be ordered directly from the publisher of
Orchideenzauber; contact Dr. Jürgen Schmidt at
E-Mail: djs@orchideenzauber.eu

The cultivation of the genus *Phragmipedium* has enjoyed great popularity for many years in Europe and North America. The author has gathered the available knowledge about this attractive genus and provided a comprehensive treatment of its species and many of its hybrids. The book also includes representatives of the genera *Mexipedium* and *Selenipedium*.

The author, a specialist in the genera *Paphiopedilum*, *Phragmipedium* and *Phalaenopsis*, has published many articles about these genera in various journals around the world and described many new species, such as *Phragmipedium dalessandroi*, *Phrag. richteri*, *Phrag. christiansenianum*, *Phrag. tetzlaffianum*, and *Phrag. brasiliense*, as well as natural hybrids *Phrag. ×colombianum*, *Phrag. ×merinoi*, *Phrag. ×pfitzerianum* and *Phrag. ×roethianum*.

The book begins with a history of the genus *Phragmipedium* dating back to 1831. General information on the genus for successful cultivation and an overview of today's nomenclature are followed by description of the plant and flower of each species, along with its variations. The geographical distribution of each species, explanation of its structure, and chromosome numbers are documented. The section for each species contains information on its distribution, location, climate, culture, blooming time, history and etymology. The author explains possible reasons for confusion in the taxonomy of this genus. A large number of pictures document the variability of each species as well as natural hybrids – some of which have never been seen before. All varieties and forma are detailed. Many hybrids are also shown. In the last part of the book the genera *Selenipedium* and *Mexipedium* are described and illustrated.

Although the book is written in German, the enormous number of pictures of this genus in one publication would make it a valuable addition to slipper lovers' libraries.

By Judith Rapacz-Hasler & Barbara Tisherman

UPCOMING EVENTS

21st WORLD ORCHID CONFERENCE

Johannesburg, South Africa, September 10-14, 2014
www.woc21.org or follow on Facebook
A Slipper Forum is being planned
Contact: Faan Kruger - info@paphiopedilum.co.za

FALL AOS MEMBERS MEETING

October 29- November 2, 2014
The 16th International Slipper Orchid Symposium will be held in conjunction with the American Orchid Society Fall Members Meeting
Location: Sheraton Orlando North,
600 N. Lake Destiny Road, Maitland, FL 32751
www.slippersymposium.com/

FALL MID-AMERICA ORCHID CONGRESS

Nov. 7-9, 2014; Central Ohio Orchid Society
Franklin Park Conservatory, Columbus, Ohio
Guest speakers include: Alexej Popov, of Popov Orchids, Germany; Andrea Niessen, of Orquideas del Valle, Colombia; and
Patricia Harding from Lebanon, OR.
www.midamericanorchids.org/meetings/

59th PAPHIOPEDILUM GUILD AND 1st WORLD SLIPPER ORCHID CONFERENCE

Saturday & Sunday, January 17 – 18, 2015
Sponsored by Orchid Digest
Expert Lectures - Paph/Phrag Displays – Orchid Sales
Hilo Hawaiian Hotel, 71 Banyan Drive, Hilo, Hawaii
Tickets: \$95 - Check the website for the latest
Information: www.orchiddigest.org

PAPHIOPEDILUM FORUM

Saturday, February 14, 2015
Washington, D.C. area
Terry Root from the Orchid Zone, CA; Theresa Hill, Hillsvie Orchids, OR; and Paul Sheetz, Just Pat Orchids, PA, are confirmed speakers.

SPRING MID-AMERICA ORCHID CONGRESS

March 20-22, 2015; Marriott Cool Springs Hotel,
Franklin, TN (hosted by the Orchid Society of
Middle Tennessee)
Guest speakers include Olaf Gruss, Germany.
www.midamericanorchids.org/meetings/

Supporting Members

In each issue of our newsletter we recognize and thank our Supporting Members (individuals and businesses) whose additional dues make it possible for us to carry out our mission of preservation and education about all lady-slipper orchids. If you are interested in becoming a Supporting Member, please contact Arlene Ricker at aricker@verizon.net. We encourage our members to support these businesses. From our website you may contact them directly.

Michael L. Ault, M.D.
 Jay Ciezki
 Jason Gebbia
 Ron L. Giles
 Steve Helbling
 Doug and Beth Martin
 Nancy Meares
 Paul Oksnee
 Judith Rapacz-Hasler
 Margaret Rente
 Arlene Ricker
 Clark T. Riley
 Jonathon M. Riley
 Barbara Tisherman
 Charles A. Walker
 Scott Ware
 Brian White
 Herb Windom
 Linda Wright & Jeff Morris
 Central Iowa OS, c/o Roger Brannian
 Deerwood Orchids, Ross & Julie Hella
 Eric Young Foundation, Eric Young
 First Rays LLC, Ray Barkalow
 Fishing Creek Orchids, Stephen Male
 Gunzenhauser Orchids, Ernst Gunzenhauser
 Hilo Orchid Farm, Inc., James Fang
 Hillview Gardens, Theresa Hill
 Hilltop Orchids, Dick & Sandy Wells
 Lehua Orchids, Graham Wood
 Little Brook Orchids, John F. Pyfer, Jr.
 Majesty Orchids, Grace Lo & Chu Yung Tsung
 Marlow's Orchids, Jim Marlow
 Marriott Orchids, Hadley & Deborah Cash
 Newman's Nursery Inc., Jeffrey Newman
 New Orleans OS, c/o Donna Strange
 Orchidaceae, Inc., Mark Srull & Joan Bateman
 Orchid Inn, Ltd., Sam Tsui

ORCHIDbabies, LLC, Earl and Phyllis Bailey
 Orchids, Ltd., Jerry and Yoko Fischer
 Orchidview Orchids, H.P. Norton
 Orosi Valley Orchids, Russell M. Tyler, Ph.D.,L.P.
 Phipps Conservatory & Botanical Gardens, Georgia Wahl
 Piping Rock Orchids, Glen F. Decker
 R.K. Gems Orchids & Tropicals, Rod Knowles
 Ravensision Photography, Jay Norris & Max Wilson
 Richard Shafer's Orchids, Richard Shafer
 Seagrove Orchids LLC, Linda Thorne
 Salventi-Purviance, John Salventi
 Slippertalk.com, c/o Heather Gaillot
 Sunset Valley Orchids, Fred Clarke
 The Tiny Jungle, Chris Mende
 Windy Hill Gardens, Marilyn & Brian LeDoux
 Woodstream Orchids, William Goldner &
 Lynn Evans-Goldner
 Zephyrus Orchids, John & Nicole Doherty

SOA DIRECTORY

President: Barbara Tisherman, Pittsburgh, PA;
 (412) 683-0207; tishermanb@gmail.com

Vice President: Lois Dauelsberg, Woodland Park, CO;
 (719) 687-2528, loisdauels@aol.com

Secretary: Alexa Noel, Cincinnati, OH
 513-469-8767; ahn6175@yahoo.com

Treasurer: Linda Thorne, Seagrove, NC;
 (336) 879-6677; orchidlinda@rtmc.net

Director: Fred Clarke, Vista, CA
 760-639-6255; fred.clarke@att.net

Director: Glen F. Decker, Galway, NY;
 (518) 882-9088; pipingrock@aol.com

Director: Steve Helbling, Cincinnati, OH
 513-321-3702; orchiddevil@aol.com

Webmaster: Rod Knowles, Brainerd, MN;
 (218) 829-4840; rkgems@brainerd.net

Journal Editor: Judith Rapacz-Hasler, Fort Myers, FL;
 (239) 223-5540; jorapacz@wisc.edu

Membership Secretary: Arlene Ricker, Canonsburg, PA;
 (724) 873-1904; aricker@verizon.net

Do you love Cymbidiums and Paphiopedilums?

The Cymbidium Society of America invites you to join

Membership includes six issues of the colorful CSA Journal magazine

Annual dues: US\$35.00 (includes overseas surface postage) add \$20.00 for overseas airmail

[VISA, MASTERCARD, JCB or checks in US funds only]

c/o Clara Moura, Membership Secretary, 1742 Grove Way, Castro Valley, CA 94546